

AD-A033 369

ARMY MEDICAL RESEARCH INST OF INFECTIOUS DISEASES FR--ETC F/G 6/5
ALTERATIONS OF THE NASAL MUCOSA OF CADAVERS OF EPIDEMIC ENCEPHALITIS(U)
DEC 76 Y WATANABE

UNCLASSIFIED

USAMRIID-MUL-0528

NL


1 OF 1
ADAO33369



END

DATE
FILMED

2-77



ADA033369

AD

14 Q USAMRIID-MUL-#528

12 b.s.

TRANSLATION NO.: MUL 0528

6 TITLE: Alterations of the Nasal Mucosa of Cadavers of Epidemic Encephalitis

10 AUTHOR(S): Watanabe, Yochio/Watanabe

11 13 Dec 76

12 4P.

REFERENCE: Tr. Soc-path. Jap. 30:580-2, 1940

DISTRIBUTION STATEMENT

Approved for public release;
distribution unlimited

D D C
DISTRIBUTED
DEC 16 1976
F

U. S. ARMY MEDICAL RESEARCH INSTITUTE OF INFECTIOUS DISEASES

Fort Derrick, Frederick, Maryland 21701

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

1473

✓405039
4

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM												
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER												
4. TITLE (and Subtitle) Alterations of the nasal mucosa of cadavers of epidemic encephalitis		5. TYPE OF REPORT & PERIOD COVERED Translation												
7. AUTHOR(s) Watanabe, Y.		6. PERFORMING ORG. REPORT NUMBER MUL 0528												
9. PERFORMING ORGANIZATION NAME AND ADDRESS Tr. Soc-path Jap. 30:580-2, 1940		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS												
11. CONTROLLING OFFICE NAME AND ADDRESS USAMRIID Library Fort Detrick, Frederick, Md. 21701		12. REPORT DATE 13 December 1976												
14. MONITORING AGENCY NAME & ADDRESS(if different from Controlling Office)		13. NUMBER OF PAGES 2												
		15. SECURITY CLASS. (of this report) Unclassified												
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE												
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release: distribution unlimited.														
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)														
18. SUPPLEMENTARY NOTES														
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Encephalitis Pathohistology St. Louis Virus		<table border="1"><tr><td>100</td><td>White Section</td><td><input checked="" type="checkbox"/></td></tr><tr><td>200</td><td>Buff Section</td><td><input type="checkbox"/></td></tr><tr><td colspan="3">UNARMED</td></tr><tr><td colspan="3">JUSTIFICATION</td></tr></table>	100	White Section	<input checked="" type="checkbox"/>	200	Buff Section	<input type="checkbox"/>	UNARMED			JUSTIFICATION		
100	White Section	<input checked="" type="checkbox"/>												
200	Buff Section	<input type="checkbox"/>												
UNARMED														
JUSTIFICATION														
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		<table border="1"><tr><td>BY</td><td colspan="2">DISTRIBUTION/AVAILABILITY CODES</td></tr><tr><td>DIST.</td><td>AVAIL.</td><td>SP/STL</td></tr><tr><td>A</td><td></td><td></td></tr></table>	BY	DISTRIBUTION/AVAILABILITY CODES		DIST.	AVAIL.	SP/STL	A					
BY	DISTRIBUTION/AVAILABILITY CODES													
DIST.	AVAIL.	SP/STL												
A														

AD

Tr soc-path jap
30:580-2, 1940

Alterations of the nasal mucosa of cadavers of
epidemic encephalitis

by

Yochio Watanabe

Institute of Pathology of Keio-Gijuku University, Director:
Prof. R. Kawamura.

Our Director, Prof. R. Kawamura, gave a report in 1936 on the pathohistologic findings in the nasal mucosa of the mouse infected with encephalitis by means of instillation of Niigata and St. Louis type virus.

In the summer of 1938 and 1939 I carried out the examination of the nasal mucosa of the respiratory and vestibular region in the course of 7 autopsies (3 adults and 4 infants who had been sick for 3-15 days) of cases of Japanese epidemic encephalitis that had broken out in Tokyo. The following findings were obtained.

In the vestibular region, aside from hyperemia and edema, there were no unusual alterations. In the respiratory region the main alteration which was encountered was gelatinization of the epithelia and nasal glands (fig. 1) and cellular infiltration, (Fig. 2), which revealed a picture of acute, catarrhal inflammation accompanied by hyperemia and edema.

MUL 0528

The cellular infiltration was observed in the area of the subepithelial tissue or in the vicinity of the glandular tissue, or diffusely in the interglandular tissue. The cellular elements were often large lymphoid cells, mixed with lymphocytes and leukocytes with lobular nuclei. Very often stratified calcium deposits could be seen in the nasal gland which stained dark with hematoxylin. In one case isolated endothelial giant cells were found in the tissue.

In brief, according to the pathohistologic study, all the cases examined by me revealed a picture of slight acute catarrhal rhinitis.

Figures

1. Lower concha. 10 year old male. 5 days of disease.
2. Lower concha. 14 year old female. 9 days of disease.

FIGURES NOT INCLUDED